

Emulsifiable Concentrate (EC)

(chemical group: Aryloxyphenoxyalkanoic Acids)

ELEGANT 5 EC

quizalofop-p-ethyl 5% w/v

Systemic post-emergence pesticide for control of annual and perennial grasses

Field of application: APPLES (Apple), Kernels (Peaches), BERRIES AND SMALL FRUITS (Strawberry), VEGETABLES (Carrot, Onions, Garlic, Tomato, Pepper, Melon, Zucchini, Cabbage, Cabbage, Peas), LEGUMES (Lentils, Beans, Chickpeas), OIL SEEDS (Soybeans, Sunflower, Cotton), POTATOES, INDUSTRIAL PLANTS (Sugar beets, tobacco, Alfalfa), CALL.

Maximum number of applications: One (1).

Method of application: Spray the weed foliage with 30-40 liters of water per acre. Spray at a pressure of less than 45 PSI (2-3 atm). Use a broom injector.

How to prepare spray liquid: Fill the spray can 3/4 full with water. Add the required amount of preparation to the spray can while stirring. Fill the sprayer with the remaining water while continuing to stir.

Elements of phytotoxicity, susceptibility of varieties and any other side effects on plants or their products: It is phytotoxic to cereals and non-robust broadleaf crops.

Compatibility: Combination with broadleaf herbicides may reduce the effectiveness of the product and should be avoided.

Last treatment before harvest: Pumpkin: 7 days. Alfalfa, Lettuce, Spinach, Carrot, Lentils, Cowpeas, Chickpeas: 21 days. Sunflower, Tobacco, Apple, Peach, Potato, Bean, Strawberry: 35 days. Radishes, Peas, Garlic: 49 days. Tomato, Cabbage, Melon, Pepper: 56 days. Sugar beet, Soybeans: 60 days. Peas, Onion, Celery: 70 days. Cotton: 80 days.

Temporal stability of the product-Storage conditions: In its original inviolable packaging, in a dry, cool and well-ventilated place, away from sources of ignition and direct exposure to sunlight. It remains stable for 2 years from the date of manufacture.

Packages: 250ml / 500ml / 1LT / 3LT

THE INFORMATION WRITTEN IN THIS FORM HAS AN INFORMATIONAL CHARACTER AND DOES NOT SUBSTITUTE IN ANY CIRCUMSTANCES.

PLANT PROTECTION PRODUCTS ARE INTENDED FOR USE BY PROFESSIONAL USERS.

INSTRUCTIONS FOR USE AND PRECAUTIONS, WRITTEN ON THE LABEL MUST BE OBSERVED.

FOR MORE INFORMATION CONTACT THE TECHNICAL DEPARTMENT OF OUR COMPANY OR CONSULT THE LOCAL AGRONOMISTS.

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System and EN ISO 14001:2015 Environmental Management System



103

10/2025



www.nitrofarm.gr



info@nitrofarm.gr



Free-phone: 800 11 820 820
+30 2310 553354-7,



Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —



ELEGANT 5 EC

SCOPE & DOSAGE

Field of application: APPLES (Apple), POME FRUIT (Peach), BERRIES AND SMALL FRUITS (Strawberry), VEGETABLES (Carrot, Onion, Garlic, Tomato, Pepper, Melon, Zucchini, Cabbage, Lettuce, Spinach, Radishes, Celery, Beans, Peas, Green Peas), LEGUMES (Lentils, Kidney Beans, Chickpeas), OILSEEDS (Soybeans, Sunflower, Cotton), POTATOES, INDUSTRIAL PLANTS (Sugar beet, tobacco, Alfalfa), ORNAMENTAL.

Targets-Doses: For Annual grass weeds: 100-150 ml/acre. For Perennial weeds: 200-300 ml/acre.

Spray volume: 30-40 liters of spray liquid/acre.

Application time: Post-emergence from the 2-3 leaf stage until the weeds are weeded and before the weeds are covered by the crop.

Maximum number of applications: One (1).

REMARKS:

1. WEEDS CONTROLLED: Annual grass weeds: Alopecurus spp. (foxtails), Apera spicaventi (windgrass), Bromus spp. (oats), Dactylis spp. (fingerweeds), Digitaria sanguinalis (bloodweed), Echinochloa crus-galli (sedge), Eleusine indica (eleusine), Hordeum spp. (barley), Lolium spp. (grass), Phalaris spp. (grasses), Panicum spp. (panic), Setaria spp. (setaria), Sorghum halepense (seed fescue), Triticum spp. (wheat).

Perennial weeds: Agropyron repens (grass), Agrostis stolonifera (grass), Cynodon dactylon

(grass), Paspalum spp. (paspalum), Sorghum halepense (rhizome fescue).

2. Resistant weeds: All broadleaf.

3. Small doses are applied when weeds are small.

4. Application time: Cynodon dactylon is best controlled when they are 15-20 cm tall. Sorghum halepense is best controlled when it has a stem length of 45-50 cm. Poa annua (annual grass) is controlled when it has 1-4 leaves and at a dose of 300 ml/hectare.

5. Do not spray when the weather is cold and when the crops are not robust.

6. Special precautions for herbicides (SZ phrases): ● Store separately from other pesticides, fertilizers and seeds. ● Use separate spraying equipment for herbicides.

● Do not contaminate irrigation or drinking water and neighboring sensitive crops. ● Do not spray when it is windy. Do not double spray the edges of the zones. ● Do not increase the doses. ● Do not use it on crops for which it is not recommended. After spraying, rinse the spraying equipment thoroughly with water and detergent. ● Do not use another drug 3 days before or after applying the preparation. ● Do not sow or plant cereals before 3 months have passed since its application.

