



Paraffin oil 83% w/v or 96,7% w/w
Emulsifiable Concentrate (EC)

NITROPOL 97,6

Insecticide-acaricide contact with action
at all stages: prosperous, mobile forms and eggs.

Action spectrum:

CITRUS (Lemon, Mandarin (Clementines), Orange), APPLES (Apple, Pear),
NUCLEAR (Apricot, Cherry, Peach, Plum), Olive*, Kiwi.

Method of application: Foliage sprays.

Elements of phytotoxicity, susceptibility of varieties and any other side effects on plants or their products: The preparation is not phytotoxic when used according to the recommended uses, doses and precautions.

Compatibility: ---.

Safety interval between application and human or animal access to the culture to which the preparation has been applied: When the spray liquid on the plants has dried (24 hours).

Last operation before harvest (days): For all crops, it is determined by the time of application of the preparation.

Storage conditions, time stability of the preparation: Keep in its original inviolable packaging, in a dry, cool and well-ventilated place, at a temperature > 0oC. In these conditions it is kept stable for 2 years from the indicated date of manufacture.

PACKAGING: 100ml / 200ml / 500ml / 1LT / 5LT / 10LT / 20LT

THE INFORMATION WRITTEN IN THIS FORM HAS AN INFORMATIONAL CHARACTER AND DOES NOT SUBSTITUTE IN ANY CIRCUMSTANCES.

PLANT PROTECTION PRODUCTS ARE INTENDED FOR USE BY PROFESSIONAL USERS.

INSTRUCTIONS FOR USE AND PRECAUTIONS, WRITTEN ON THE LABEL MUST BE OBSERVED.

FOR MORE INFORMATION CONTACT THE TECHNICAL DEPARTMENT OF OUR COMPANY OR CONSULT THE LOCAL AGRONOMISTS.

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System and EN ISO 14001:2015 Environmental Management System





55

12/2020

 www.nitrofarm.gr

 info@nitrofarm.gr

 Free-phone: 800 11 820 820
+30 2310 553354-7,

 Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —

NITROPOL 97.6

SCOPE & DOSAGE

Scope - Objective - Doses - Time of application:

CITRUS (Lemon, Mandarin (Clementines), Orange):

i) Red scabies, Parlatoria and other scabies with 1000-1500 ml / 100 liters spray liquid or 1000-2240 ml in 100-220 liters of spray liquid / acre. Applications before fruit color change (BBCH 00-81).

ii) Flour with 1000-1500 ml / 100 liters spray liquid or 500-2240 ml in 50-220 liters of spray liquid / acre. Application before the change of color of the fruits (BBCH 00-81) or an autumn one.

APPLES (Apple, Pear):

i) San Jose scabies and other scabies with 750-1000 ml / 100 liters spray liquid or 600-1000 ml in 80-100 liters spray liquid / acre. Application before fruit color change (BBCH 00-81).

ii) Red nightshade with 750-1000 ml / 100 liters spray liquid or 940-1500 ml in 125-150 liters of spray liquid / acre. Application before flowering (BBCH 54-56).

NUCLEAR FRUITS (Apricots, Cherries, Peaches, Plums):

i) San Jose scabies and other scabs with 750-1000 ml / 100 liters spray liquid or 600-1000 ml in 80-100 liters spray liquid / acre. Application before fruit color change (BBCH 00-81).

ii) Red and Yellow nightshade with 750-1000 ml / 100 liters spray liquid or 940-1250 ml in 125 liters of spray liquid / acre. Application before flowering (BBCH 51-53).

Olive (Minor Uses)*:

Parlatoria, Pelion, Pollinia, White scabies, Mytilomorfi scabia, Hours, Mites (*Eriophyes vitis**, *Tegolophus hassani**, *Oxyceus maxwelli**, *Eotetranychus carpini**) with 750-1000 ml utensils / 100 liters spray liquid or 600-1500 ml in 80-150 liters spray liquid / acre. Application with the appearance of the infestation, from the fall of the petals to small fruits (BBCH 67-79).

Kiwi (Use of Minor Significance)*:

San Jose Sanas, Cotton or Dispis, Metcalfa, Cotton, Yellow Tetranychus, Red Tetranychus with 750-1000 ml / 100 liters spray liquid or 600-1500 ml in 80-150 liters of spray liquid / acre. Application with the appearance of the infestation, before flowering and after harvest (BBCH 50-90).

Maximum number of applications per growing season / interval of applications in days:

CITRUS (Lemon, Mandarin (Clementines), Orange): 1-2 /> 60, APPLES (Apple, Pear), NUCLEAR (Apricot, Cherry, Peach, Damask), Olive, 1.

OBSERVATIONS:

1. The range of doses and the volume of spray liquid depends on the height of the plants and the intensity of the infestation.
2. In citrus fruits the recommended dose should be combined with the appropriate volume of spray liquid so as not to exceed the maximum dose / acre.
3. When using **NITROPOL 97.6** the trees must be watered. Water one week before spraying.
4. Avoid spraying when plants are stressed, during periods of intense drought, excessive humidity and when there is frost.
5. Do not apply with the preparation, during the flowering period or when the temperature exceeds 32°C because phytotoxicity problems may occur.
6. Do not apply with the preparation for 30 days before or after the application of sulfur or calcium sulfate.
7. Do not precede or follow the application of preparations with d.o. captan, folpet and dithianon for at least 1 week.

* "The effectiveness has not been documented and no control has been done for possible negative effects on olive and kiwi crops from the use of the preparation. For these uses, the licensee is not responsible for possible failures related to the effectiveness or phytotoxicity of the use of the preparation".

56

12/2020



www.nitrofarm.gr



info@nitrofarm.gr



Free-phone: 800 11 820 820
+30 2310 553354-7,



Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —

