



Product of parallel trade (reference product: BELPROMEC 1,8 EC)
abamectin 1.8% • Emulsifiable Concentrate (EC)

Insecticide acaricide for stomach and contact, with extruding action against mobile forms of insects and mites. It causes paralysis in insects and mites resulting in their death from starvation, within 3-5 days of spraying.



Spectrum of action: Field of application - Target - Dosage of preparation, Volume of spray liquid (liters/acre) - Method and time of application - Number of applications per growing season, Application interval in days: **Strawberry (G):** i) Yellow spider mite, *Phytomemus pallidus*, 15-100 ml/acre, 50-100 ml/100 liters of spray liquid, 30-100 liters/acre. 1-2 applications every 7 days with the appearance of the first mobile forms. ii) *Liriomyza*, 15-100 ml/acre, 50-100 ml/100 liters of spray liquid, 30-100 liters/acre. 1-2 applications every 7 days with the appearance of the infestation.

Tomato (G): i) Yellow spider mite, *Eryophyidae* (*Aculops lycopersici*), *Polyphagotarsonemus latus*, 15-100 ml/ha, 50-100 ml/100 liters of spray liquid, 30-100 liters/ha. 1-3 applications every 7 days with the appearance of the first mobile forms. ii) *Liriomyza*, 15-100 ml/ha, 50-100 ml/100 liters of spray liquid, 30-100 liters/ha. 1-3 applications every 7 days with the first egg laying. iii) *Tuta absoluta*, 100 ml/ha, 100 ml/100 liters of spray liquid. liquid, 100 liters/hectare. 1-3 applications every 7 days upon the appearance of the infestation.

Eggplant (G): Yellow four-claw, 15-80 ml/str, 50-80 ml/100 liters spray. liquid, 30-100 liters/sq. 1-3 applications per 7 days with the appearance of the first mobile forms.

Pumpkin (G), Cucumber (G), Melon (G), Watermelon (G), Pumpkin (G): i) Yellow four-claw, 15-100 cc/str, 50-100 cc cm./100 liters of spray liquid, 30-100 liters/sq. 1-2 applications per 7 days with the appearance of the first mobile forms. ii) *Liriomyza*, 15-100 ml/str, 50-100 ml/100 liters spray. liquid, 30-100 liters/sq. 1-2 applications every 7 days with the first spawning.

Ornamental and floricultural (G): *Tetranychus*, 15-100 ml/str, 50-100 ml/100 Liters spray. liquid, 30-100 liters/sq. 1-3 applications per 15 days with the appearance of the first mobile forms.

NOTES: Large doses are applied when large populations are present.

Special agricultural, phytosanitary or environmental conditions under which the formulation can be used or excluded: The formulation must not be applied to greenhouse crops: Pumpkin, Cucumber, Cucumber, Melon, Watermelon, Squash, Strawberry, during the winter months (November- February).

Resistance management: To avoid the development of resistance it is recommended to alternate with plant protection products with a different mode of action. Do not use it for more than two consecutive applications.

Safety interval between application and: -sowing or planting of the protected crop: --, -sowing or planting of the following crops:--,-human or animal access to the crop to which the preparation has been applied: When the spray liquid has dried on the plants.

Method of application: Foliar spraying.

How to make spray liquid: Fill the sprayer tank halfway with water. Add the recommended amount of formulation to the spray can, stirring. Top up with the rest of the water while continuing to stir.

Cleaning sprayers: Empty the spray can completely. Make sure all traces of the preparation have been removed. Rinse the barrel and all parts of the sprayer three (3) times with clean water.

Evidence of phytotoxicity, cultivar sensitivity and any other side effects on plants or their products: It is not phytotoxic at the recommended uses and application rates.

Last operation before harvest or before placing on the market when it comes to post-harvest uses (days): Strawberries (G), Tomatoes (G), Eggplants (G), Zucchini (G), Cucumbers (G), Cucumbers (G), Melons (G), Watermelons (G), Pumpkins (G): 3.

Storage conditions, time stability of the formulation: Store in its original closed packaging, in a cool, dry and well-ventilated area, away from sources of ignition and heat. In these conditions it remains stable for two (2) years from the date of its manufacture.

THE INFORMATION WRITTEN IN THIS FORM HAS AN INFORMATIONAL CHARACTER AND DOES NOT SUBSTITUTE IN ANY CIRCUMSTANCES.

PLANT PROTECTION PRODUCTS ARE INTENDED FOR USE BY PROFESSIONAL USERS.

INSTRUCTIONS FOR USE AND PRECAUTIONS, WRITTEN ON THE LABEL MUST BE OBSERVED.

FOR MORE INFORMATION CONTACT THE TECHNICAL DEPARTMENT OF OUR COMPANY OR CONSULT THE LOCAL AGRONOMISTS.

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System and EN ISO 14001:2015 Environmental Management System

32

01/2026



www.nitrofarm.gr



info@nitrofarm.gr



Free-phone: 800 11 820 820
+30 2310 553354-7,



Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —