



NEW

WETTABLE GRANULES (WG) ● boscalid 26,7% w/w + pyraclostrobin 6,7% w/w

CHILA

Fungicide with preventive and, in some cases, therapeutic action that prevents spore germination, mycelium development and sporogenesis.

The d.o. pyraclostrobin interferes with cellular respiration at the Qo site of complex III of the respiratory chain and exhibits aplastic and delamination movement, while d.o. boscalid inhibits cellular respiration but at a different site of action (complex II) and exhibits transsystemic movement.



Field of Application - Target: Tomato (F) for *Alternaria* (*Alternaria* sp.).

Dose: 150 gr. at 50-100 liters of spray/hectare.

Method and Time of Application: Preventive applications when the conditions are favorable for the development of the diseases from the beginning of the formation of the lateral shoots until the appearance of the typical color in 70% of the fruits (BBCH 20-87).

Maximum number of applications / growing season: Up to 3 applications / 8-10 days.

Specific agricultural, phytosanitary or environmental conditions under which the formulation may be used or excluded: **Resistance management:** ● The formulation contains the active substances boscalid and pyraclostrobin, which belong to FRAC groups 7 and 11, respectively. ● It is recommended not to use the formulation for more than 2 consecutive applications. Before the 3rd application, use an active substance with a different mode of action. ● It is recommended to apply the preparation preventively or in the first stages of the development of diseases. Avoid therapeutic applications.

Safety interval between application and i) sowing or planting of the protected crop: In case of failure of the crop to which the formulation was applied, the same field can be re-sown after one (1) year, ii) **sowing or planting of the following crops:** Do not sow the next crop before one (1) year has passed since the application of the formulation, iii) **human or animal access to the crop to which the formulation has been applied:** Do not enter the sprayed area before the spray liquid has completely dried from the leaves.

Evidence of phytotoxicity, cultivar sensitivity and any other side effects on plants or their products: Not phytotoxic in the proposed crops and application rates.

Method of application: Foliage coverage spraying.

How to prepare the spray liquid: Fill the sprayer tank halfway with water. Add the required amount of the formulation to the spray can while stirring. Top up with the rest of the water while continuing to stir.

Cleaning sprayers: Empty the spray can completely. Rinse the barrel and all parts of the sprayer with water and detergent, then rinse three (3) times with clean water.

Instructions for the safe disposal of the plant protection product and packaging: The empty packaging (bottles) are rinsed under pressure with a suitable mechanism or triple rinsed (the rinse water is poured into the spray container) and then after having been previously damaged by puncturing or tearing (bottles, bags, boxes), to ensure no further use, are deposited at collection points for recycling or energy recovery.

Combinability: --.

Last operation before harvest or before placing on the market in the case of post-harvest uses (days): Tomato: 3.

Storage conditions, time stability of the formulation: The formulation remains stable for two (2) years from the date of manufacture, when stored in its original sealed packaging in a dry, cool and well-ventilated area.

THE INFORMATION WRITTEN IN THIS FORM HAS AN INFORMATIONAL CHARACTER AND DOES NOT SUBSTITUTE IN ANY CIRCUMSTANCES.

PLANT PROTECTION PRODUCTS ARE INTENDED FOR USE BY PROFESSIONAL USERS. INSTRUCTIONS FOR USE AND PRECAUTIONS, WRITTEN ON THE LABEL MUST BE OBSERVED.

PACKAGES:

100gr / 150gr / 500gr / 1Kg / 2Kg



57

01/2024

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System and EN ISO 14001:2015 Environmental Management System



www.nitrofarm.gr



info@nitrofarm.gr



Free-phone: 800 11 820 820
+30 2310 553354-7,



Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —