



POTASSIUM SULPHATE
LIQUID FERTILIZER
POTASSIUM - SULPHUR

K-S ultimate

FERTILIZER



NPK (S) 0-0-25 (42) / 0-0-20,75 (16,8)
PFC 1(C)(I)(b)(i): SIMPLE LIQUID INORGANIC MACRONUTRIENT FERTILIZER
25% potassium oxide (K₂O) (=20.75% K) water soluble
42% sulphur trioxide (SO₃) (=16.8% S) water soluble

Ingredients: Potassium thiosulfate (CAS No. 10294-66-3)
KSY1, Substances and mixtures of virgin materials.

Liquid in solution: ● Low in chlorides ● Density approximately 1.475 kg/liter.

Storage conditions: The product is stable under normal storage conditions (0-40°C).

Store in a cool and well-ventilated place.

In case of accidental release, absorb with sand or earth and collect mechanically in suitable containers.

Keep containers tightly closed and properly labeled.

Safety and environmental information:

To avoid risks to human health and the environment, follow the recommended instructions for use of this fertilizer.

POISON CENTER TEL.: 210-7793777

Do not disperse the container in the environment.

Consult the safety data sheet.

General information: Avoid mixing with oxidizing agents and strong acids.

FOR PROFESSIONAL USE ONLY.

Contact with acids releases toxic gases.

**It's not a plant
protection product**



PAGE 50

NitroFarm SA is certified with EN ISO 9001:2015 Quality Management System
and EN ISO 14001:2015 Environmental Management System

Potassium thiosulfate is suitable for all crops in the biological stages of plants, where potassium needs are increased.

General dosage: 1-2 liters / 1000m² through the irrigation system or foliar 200-400 ml / 100 liters of water. Repeat when necessary every 15-20 days.

Stone fruits, giant fruits: 1-3 applications in total. 1st application at the beginning of color change with 200-300 ml / 100 liters of water and 2nd application 10-15 days later with the same dosage. Repeat if necessary in 10 days with the same dosage.

Citrus fruits: 1st application at the beginning of fruit ripening with 200-300 ml / 100 liters of water. / 100 liters of water. 1-2 repetitions with the same dosage.

Olive: 1st application at the 'polishing' of the fruit with 200-300 ml. / 100 liters of water and repetitions every 20-25 days with 200-400 ml. / 100 liters of water.

Grapevine: 2-3 applications from the stage of perk with 200-300 ml. / 100 liters of water and every 15-20 days with the same dosage.

Vegetables, strawberry: 300-400 ml. / 100 liters of water at the beginning of the change in color of the fruits. 1-2 liters / 1000m² for applications through the irrigation system.

Potato: 1st application after the start of tuberization with 200-400 ml / 100 liters of water or with 2-3 liters / 1000m² with the irrigation system. Repeat in 10 days with the same dosage.

Carrot, garlic: Foliar sprays with 200-300 ml / 100 liters of water from the formation of the rhizome onwards.

COMPATIBILITY:

Potassium Thiosulfate is combined with all common pesticides and fertilizers. In spray solutions of Potassium Thiosulfate, the pH should not be less than 6.

It is also not advisable to apply to soils with a pH lower than 6.

In any case of unknown combination, a test should be carried out on a small area beforehand.

It is applied during the coldest hours of the day and immediately after preparing the solution.

For spraying in greenhouses, apply the smallest doses.



www.nitrofarm.gr



info@nitrofarm.gr



Free-phone: 800 11 820 820
+30 2310 553354-7,



Industrial Area of Sindos, SQ20,
B35-55, 57022, Thessaloniki

NitroFarm

— since 1976 —